





Obesity Facts

**32nd European Congress on Obesity
(ECO 2025)**

Malaga, Spain, May 11–14, 2025

Abstracts

EASO



Epigenetic signatures in saliva of offspring from mothers with Gestational Diabetes: longitudinal associations with early growth

Pino Ariza, L.¹; Linares Pineda, T.²; Lendínez Jurado, A.⁴; Piserra López, A.⁵; Suárez Arana, M.⁶; Pozo, M.²; Molina Vega, M.³; Picón César, M. J.³; Martínez Montoro, J. I.³; Morcillo, S.³

¹Universidad de Málaga Campus de Teatinos s/n Málaga Spain

²Instituto de Investigación Biomédica de Málaga y Plataforma en Nanomedicina-IBIMA Plataforma BIONAND

³Servicio de Endocrinología y Nutrición Hospital Universitario Virgen de la Victoria, Instituto de Investigación Biomédica de Málaga y Plataforma en Nanomedicina-IBIMA Plataforma BIONAND Málaga Spain, Centro de Investigación Biomédica en Red de la Fisiopatología de la Obesidad y la Nutrición (CIBEROBN) Instituto de Salud Carlos III Madrid Spain

⁴Departamento de Endocrinología Pediátrica Hospital Regional Universitario de Málaga Spain, Universidad de Málaga Spain

⁵Departamento de Cardiología Hospital Universitario Virgen de la Victoria Málaga Spain

⁶Departamento de Obstetricia y Ginecología Hospital Regional Universitario de Málaga Spain

Introduction: The prevalence of obesity and type 2 diabetes mellitus (T2DM) is rising globally, particularly among children exposed to adverse intrauterine environments, such as those associated with gestational diabetes mellitus (GDM). Epigenetic modifications, specifically DNA methylation, have emerged as mechanisms by which early environmental exposures can predispose offspring to metabolic diseases. This study aimed to investigate DNA methylation differences in children born to mothers with GDM compared to those born to mothers without GDM, using saliva samples, and to assess the association of these epigenetic patterns with early growth measurements.

Methods: This study analyzed saliva DNA methylation patterns in 30 children (15 born to mothers with GDM and 15 to mothers without GDM) from the EPIDG cohort. Samples were collected at two time points: 8–10 weeks postpartum and at one year of age. Epigenome-wide analysis of over 850,000 CpG sites was conducted using the Illumina Methylation EPIC Bead Chip. Differential methylation positions (DMPs) were identified with the limma package, using a significance threshold of $p < 0.01$ and $\Delta\beta \geq 5\%$. Correlation analysis examined associations between methylation and growth variables (weight, height, BMI and annual growth) using Spearman tests.

Results: We identified 6,968 DMPs at the postpartum stage and 5,132 after one year, with 50 sites remaining differentially methylated over time, 16 of which maintained consistent methylation directionality. Functional analysis linked several of these DMPs to genes involved in inflammation and metabolic processes, including CYTH3 and FARP2, both implicated in growth and metabolic pathways. Significant correlations were found between specific CpG sites and growth-related variables such as weight, head circumference, height, and BMI.

Conclusions: This study's longitudinal design reveals stable DNA methylation patterns in saliva samples that differentiate children born to mothers with GDM compared to those born to mothers without GDM across the first year of life, highlighting the feasibility of saliva as a minimally invasive biomarker source. The persistence of these epigenetic signatures underscores their potential as early indicators of metabolic risk, offering valuable insights into the long-term impact of GDM during pregnancy on children's health and development. Although the use of saliva offers a practical and non-invasive tool for paediatric epigenetic research, further studies are necessary to validate these findings in larger populations.

*First and second authors contributed equally to this work.

Co-Creation of Digital Health Interventions with Children: Initial Results of the Multi-Country Workshop in the BIO-STREAMS Project

Mrakar, I.¹; Smrke, U.¹; Rehberger, A.¹; Charmandari, E.²; Kassari, P.²; Karachaliou, A.³; Bigec, M.⁴; Bander, A.⁵; Langlet, B.⁶; Gernay, C.⁷; Jiang, Z.⁸; Martello, G.⁹; Hansov, S.¹⁰; Panagiotidou, M.¹¹; Kakkos, I.¹²; Anastasiou, A.¹²; Salamon, E.¹³; Doran, R.¹⁴; Anjos, S.¹⁵; Lambrou, D.¹⁶; Ciudin, A.¹⁷; Georga, E. I. I.¹⁸; Fotiadis, D. I. I.¹⁹; Prasinos, M.²⁰

¹University of Maribor, Faculty of Electrical Engineering and Computer Science

²Center for the Prevention and Management of Overweight and Obesity in Childhood and Adolescence, Division of Endocrinology, Metabolism and Diabetes, First Department of Pediatrics, School of Medicine, National and Kapodistrian University of Athens

³Department of Nutrition and Dietetics, School of Health Sciences and Education, Harokopio University of Athens

⁴University Division of Pediatrics, University Medical Centre Maribor

⁵Danish Committee for Health Education

⁶Department of Medicine, Huddinge, Karolinska Institutet

⁷Department of Pediatrics, CHU of Liège

⁸Department of Information System Management, CHU of Liège

⁹MARTEL GmbH

¹⁰Blocks Health and Social Care

¹¹AINIGMA Technologies

¹²Institute of Communications and Computer Systems

¹³European School Heads Association

¹⁴NUCLIO

¹⁵ICS

¹⁶Penteli Children's Hospital

¹⁷VHIR - Vall d'Hebron Institute of Research

¹⁸Unit of Medical Technology and Intelligent Information Systems, Dept. of Materials Science and Engineering, University of Ioannina

¹⁹Biomedical Research Institute, Foundation for Research and Technology-Hellas, Ioannina

²⁰Telematic Medical Applications

Background: Digital health interventions show promise in preventing childhood obesity, yet their effectiveness depends heavily on understanding and incorporating users' values and preferences. Limited research exists on value-sensitive design approaches incorporating children's perspectives across different European contexts.

Objective: To identify key values, preferences, and requirements of children for digital health interventions. These findings will directly inform the design of BIO-STREAMS digital solutions and guide the design and implementation of BIO-STREAMS prospective studies aimed at childhood obesity prevention.

Methods: Structured workshops with children (aged 8 to 18) in educational and clinical settings, in Belgium, Bulgaria, Denmark, Greece, Netherlands, Slovenia, Spain, Sweden, and Portugal. We are using participatory design methods including presentations, demonstrations, mock-ups, group discussions, and interactive activities to explore children's perspectives on healthy lifestyles, digital learning preferences, and engagement requirements. Data is analyzed using thematic analysis approach.

Results: Workshops included 23 children from clinical (still ongoing) and 181 from educational settings. Initial findings show setting-specific differences: children in clinical settings prioritized health tracking and structured approaches, while those in educational settings favored social interaction. Age-specific patterns emerged from play-based learning in younger children (<8y) to complex digital engagement in adolescents (14-18y). Overall, key requirements include age-adaptive interfaces. These should progress from visual/play-based to social/autonomous features, balancing individual health tracking with social engagement. All interventions and interfaces should incorporate interactive elements, cultural sensitivity, achievement systems, flexible parental controls, and privacy features.

Conclusions: Initial findings emphasize the need for age-specific customization in digital health interventions and balance between digital and in-person interaction. Effective interventions should include social features, cultural sensitivity, and age-appropriate achievement systems with contextual flexibility. These insights can inform the design of more engaging and sustainable childhood obesity prevention programs. The results of workshops have already been used to inform the design of BIO-STREAMS digital interventions and can be used in the future for more engaging, long-term sustainable and more effective childhood obesity prevention programs.

Funding: This work has been funded by the European Union Horizon Europe Research and Innovation Programme project BIO-STREAMS (grant number 101080718), Swiss State Secretariat for Education, Research and Innovation (SERI), and the Slovenian Research and Innovation Agency, Advanced Methods of Interaction in Telecommunication research programme (grant number P2-0069). The content of this work does not reflect the official position of the European Union or any other institution. The information and views expressed are the sole responsibility of the authors.

LBP3.06

Mediterranean diet adherence, sleep adequacy and obesity prevalence among students in remote areas of Greece: A cross-sectional study

Maraki, M.¹; Kanta, A.²; Vasiliou, T.³; Kyritsi, E.³; Koulouvaris, P.⁴

¹Section of Sport Medicine and Biology of Exercise, School of Physical Education and Sport Science, National and Kapodistrian University of Athens, Athens, Greece

²Department of Nutrition and Dietetics Sciences, Hellenic Mediterranean University, Sitia, Greece

³Mobile Medical Units, Stavros Niarchos Foundation, Greece

⁴1st Department of Orthopedic Surgery, National and Kapodistrian University of Athens, Athens, Greece

Introduction: Childhood obesity is a multifactorial disease and a significant global public health concern. Lifestyle factors, including dietary habits and sleep patterns, have been linked to weight status. Additionally, limited access to healthcare services in remote areas has been associated with poorer health outcomes, including higher obesity prevalence. This study aimed to assess the prevalence of overweight and obesity in a large sample of children living in remote rural areas of Greece and to examine the association between adherence to the Mediterranean Diet (MD), sleep adequacy and weight status.

Methods: This cross-sectional, population-based observational study included children aged 8 to 18 years, recruited from primary and secondary public schools in 29 remote areas of Greece, both island and mainland. Anthropometric measurements were recorded, and adherence to the MD was assessed using the KIDMED questionnaire. Sleep duration was evaluated based on self-reported wake-up and bedtime on weekdays and weekends. Multivariable logistic regression was employed to assess associations between diet quality, sleep adequacy, and the likelihood of overweight and obesity.

Results: A total of 938 students participated in the study, including 458 boys (48.8%) and 480 girls (51.2%). Among them, 52.5% resided in remote mainland areas, while 47.5% lived on remote islands. The prevalence of overweight, obesity, and central obesity was 24.7%, 14.1%, and 36.9%, respectively. In terms of MD adherence, 15.4% of students had low adherence, 50.1% had moderate adherence, and 34.6% had high adherence. Regarding sleep duration, 15.2% of students were classified as short sleepers and 5.2% as long sleepers. No significant differences were observed between boys and girls.

Students with low MD adherence were 2.5 times more likely to be overweight or obese (OR 2.52, 95% CI: 1.66–3.83, $p < 0.001$) and 1.6 times more likely to have central obesity (OR 1.63, 95% CI: 1.06–2.53, $p = 0.021$)

compared to those with high MD adherence. Furthermore, students with low MD adherence had 1.8 times the likelihood of being overweight or obese (OR 1.83, 95% CI: 1.24–2.71, $p = 0.003$) compared to those with moderate adherence. Additionally, students with moderate MD adherence were 37% more likely to be overweight or obese (OR 1.37, 95% CI: 1.01–1.88, $p = 0.044$) than those with high adherence.

Short sleepers were 2.6 times more likely to be overweight or obese (OR 2.63, 95% CI: 1.38–5.01, $p = 0.003$) compared to students with adequate sleep duration. Notably, students with low MD adherence were three times more likely to be short sleepers (OR 3.22, 95% CI: 1.26–8.2, $p = 0.014$) compared to those with high adherence.

Conclusion: Consistent with previous studies in urban populations, our findings suggest that among students living in remote mainland and island areas, higher adherence to the MD is associated with a lower risk of overweight and obesity, while short sleep duration is linked to a higher risk. These results highlight the importance of promoting the MD and adequate sleep in remote areas as key strategies for combating obesity.

LBP3.07

Harms Reporting in Pharmacological Randomized Controlled Trials in Children and Adolescents with Obesity: A Systematic Review

Pietrasik, M.¹; Kawish, M.¹; Silva, H.¹; Sacco, S.²; Carrière, K.¹; Wahi, G.³; Ball, G.⁴; Johnston, B.⁵; Merdad, R.⁶; Jebeile, H.⁷; Sherifali, D.⁸; Fitzpatrick Lewis, D.⁸; Usman Ali, M.⁹; Morrison, K. M.³; Alberga, A.¹

¹Department of Health, Kinesiology, And Applied Physiology, Concordia University, Montreal, Quebec, Canada

²Department of Psychology, Concordia University, Montreal, Quebec, Canada

³Department of Pediatrics, McMaster University, Hamilton, Ontario, Canada

⁴Department of Pediatrics, Faculty of Medicine and Dentistry, College of Health Sciences, University of Alberta, Edmonton, Alberta, Canada

⁵Department of Nutrition, Texas A&M University, College of Agriculture and Life Sciences, College Station, Texas, USA

⁶King Abdulaziz University, Jeddah, Saudi Arabia

⁷The University of Sydney, Children's Hospital Westmead Clinical School, Westmead, Australia

⁸School of Nursing, Faculty of Health Sciences, McMaster University, Hamilton, ON, Canada

⁹Department of Health Research Methods, Evidence and Impact, Faculty of Health Sciences, McMaster University, Hamilton, Ontario, Canada

Introduction: Medications (e.g., GLP-1 receptor agonists, metformin) are increasingly being used in pediatric obesity management. This systematic review aimed to (i) explore the extent of harms reporting in pharmacological randomized controlled trials (RCTs) for managing obesity in children and adolescents and (ii) identify the types of harms (i.e., physical vs. psychosocial) reported.

Methods: A database search was conducted to identify RCTs of pharmacological interventions for children and adolescents (median age <18 years) with obesity in MEDLINE, Embase, and Cochrane Library published up to November 2022. Two reviewers independently screened articles and extracted data on harms reporting using the Consolidated Standards of Reporting Trials (CONSORT) Harms 2022 Statement guidance for reporting harms in RCTs and other detailed questions about harms reporting created by our research team.

Results: 35 trials (published from 2001 to 2022) were included, and 30 reported harms. There were no studies that reported on all CONSORT Harms items. No trials reported a definition of “harms” or how harms were measured (e.g., based on a discrete definition, visual analogue scale). Only two studies (6%) reported adhering to CONSORT Harm-reporting standards, five (17%) reported open accessibility of harms data, and seven (23%) reported the statistical analytic method used to assess harms. Of the 30 studies that reported harms, 29 studies (97%) reported multiple